




Greenland Rift Valley Surface of Earth (crust), Floating on a layer of denser material. North America Alfred Wegener 1912-1915 observations Africa and South America fit Hypothesis: Africa Continental drift Pigure 6-5
Discovering the Universe, Eighth Edition

© 2008 W.H. Freeman and Company





9 10





11 12







13 14







15 16







17 18







19 20






21 22







23 24





25 26






27 28






This photograph, made with a microscope, shows tiny microcraters less than 1 mm across on a piece of Moon rock.





29 30




Using a laser mounted on the *Lunar Reconnaissance Orbiter*, this detailed image of the lunar far side was made in 2010. Going by the colors of the rainbow, violet indicates lowest terrain, while red indicates highest.


31 32






33 34






35 36




Seismic experiments revealed that the main regions of the Moon's interior mimic those of Earth, but in different proportions.


Water ice may exist in the polar craters, where the energy received from the Sun is insufficient to melt it.

37 38






39 40



The Moon probably formed from debris cast into space when a huge planetesimal struck the proto-Earth


The collisional-ejection theory holds that the proto-Earth was struck by a Mars-sized protoplanet and that debris from this collision coalesced to form the Moon
This theory successfully explains most properties of the Moon
The Moon was molten in its early stages, and the anorthositic crust solidified from low-density magma that floated to the lunar surface
The mare basins were created later by the impact of planetesimals and filled with lava from the lunar interior

41 42




As the Moon's interior shrank, the surface settled irregularly, creating long lines of diffs called scarps.

43 44





45 46




Why do we see only one side of the moon from earth?


Tidal forces cause the moon to be in synchronous rotation.







49 50






51 52







53 54



The central indentation in this Messenger image from 2009 is believed to be a caldera (sunken vent) of an explosive volcano on Mercury. It is unlikely to be an impact crater, as it completely lacks a raised crater wall.

55 56





57 58

Mercury has an iron core and a surprising magnetic field

- Most iron-rich (relative to mass) planet in the solar system with a core that is 75% of the diameter
- The earth's core is 55% of its diameter and the moon's core is 20% of its diameter
- Among highest density for the planets
- Weak magnetic field indicating part of the core is liquid
- Magnetic field causes a magnetosphere similar to Earth's but weaker

Mantle

Earth

In the entire planet, or 17% of its volume...

In the entire planet, or 17% of its volume...


Mantle


Mercury

Mantle


Mercury

59 60






61 62






63 64







65 66




Venus has a hot, dense atmosphere and highly reflective corrosive cloud layers Temperature (°C) Spacecraft measurements 100-100 100 200 300 400 reveal that 96.5% of the 0.0001 Venusian atmosphere is carbon dioxide 0.001 80 Most of the balance of the atmosphere is nitrogen. 0.01 0.1 Upper cloud layer Pressure (atm) Venus's clouds consist of £ 60 Middle cloud layer Lower cloud layer droplets of concentrated sulfuric acid. Haze layer 40 The surface pressure on Venus is 90 atm, and the surface temperature is 20 460° C Both temperature and pressure decrease as J90 200 300 400 500 600 700 altitude increases Temperature (K)


67 68







09







71 72







73 74






75 76





77 78





(a) Mars from the Hubble Space Telescope

For reasons that are not understood, the chemical composition of ancient Martian lava is different from that of more recent lava


Mars has no planet wide magnetic field at present but may have had one in the ancient past


79 80







81 82







83 84







85






87 88





89 90




Olympus Mons,


Disconvinta Eighth delicion

0 2008 W/A Freema and Congary

the larges volcano on Mars and in the solar system

91 92







93 94





95 96







97 98






99 100






101 102




The two Martian moons resemble asteroids



- Mars has two small, football-shaped satellites that move in orbits close to the surface of the planet
- They may be captured asteroids or may have formed in orbit around Mars out of solar system debris

103 104



Why is there no H_2 in the atmosphere of the inner planets?

 $\begin{array}{llll} E_k = 1/2 m v^2 & J & \text{Kinetic energy due to motion with velocity, v} \\ E_k = 3/2 k T & J & \text{Thermal kinetic energy of gas, atoms or molecules} \\ v = (3kT/m)^{1/2} & m/s & \text{Average speed of a gas, atom or molecule} \\ k = 1.38 x 10^{-23} & J/K & \text{Boltzmann's constant} \\ m = \mathcal{M} \ x \ amu & kg & \text{mass of atom or molecule} \\ \mathcal{M} & \text{mass number} \\ amu = 1.66 \ x \ 10^{-27} \ kg & \text{atomic mass unit} \end{array}$

A planet or moon can retain a gas if the escape speed is at least 6 times greater than the average velocity of the gas.

105 106

What is the escape velocity for Mars?

 v^2_{escape} = 2GM/r M = 6.418x10²³ kg (mass of Mars) r=3397 km (radius of Mars)

 $v^2_{\rm escape} = 2 \ x \ 6.673 x 10^{-11} \ x \ 6.418 x 10^{23} \ / (3.397 \ x \ 10^6) = 25.214 \ x \ 10^6 \ (m/s)^2 \ V_{\rm escape} = 5021 \ m/s = 5.021 \ km/s$

Why is there no H_2 in the atmosphere of the inner planets?

 $E_k = 1/2 m v^2$ Kinetic energy due to motion with velocity, v $E_k=3/2kT$ Thermal kinetic energy of gas, atoms or molecules $v = (3kT/m)^{1/2}$ m/s Average speed of a gas, atom or molecule k=1.38x10⁻²³ J/K Boltzmann's constant mass of atom or molecule $m = \mathcal{M} \times amu$ kg mass number amu= 1.66 x 10⁻²⁷ kg atomic mass unit

Example for Mars: T=220K, H_2 : \mathcal{M} =2, m= 2x1.66x10⁻²⁷ kg v=[(3x1.38x10⁻²³x220/(2x1.66x10⁻²⁷)]^{1/2}=1656 m/s

A planet or moon can retain a gas if the escape speed is at least 6 times greater than the average velocity of the gas.

 V_{esc} =5.02 km/s =3.0 x v \rightarrow H₂ cannot be retained by Mars.

107 108