



Biophysics I (BPHS 4080)


Instructors: Prof. Christopher Bergevin (cberge@yorku.ca)

Website: <http://www.yorku.ca/cberge/4080W2018.html>

Water flow through semipermeable membrane



$$\Phi_V = \mathcal{L}_V [(p^i - \pi^i) - (p^o - \pi^o)]$$




Kinetic Volume Changes in Cell Volume

$$\Phi_V = \mathcal{L}_V [(p^i - \pi^i) - (p^o - \pi^o)]$$

assume no hydraulic pressure


Primary osmotic responses of cells



→ outward flux must equal reduction
in intracellular water volume
(per unit area of membrane)



$$\frac{d\mathcal{V}^i(t)}{dt} = -A(t)RT\mathcal{L}_V(C_\Sigma^o(t) - C_\Sigma^i(t))$$

Primary osmotic responses of cells




Equilibrium Solution

$$v_C(\infty) = v_C' + \frac{N_{\Sigma}^i}{C_{\Sigma}^o} \rightarrow \text{perfect osmometer}$$




→ Cells can actually behave in this way!

Osmotic Response Dynamics: Egg Cell




$$\begin{aligned}
 \frac{d\mathcal{V}^i(t)}{dt} &= 4\pi r^2(t) \frac{dr(t)}{dt} \\
 &= 4\pi r^2(t) RT\mathcal{L}_V \left[\frac{N_\Sigma^i}{\frac{4}{3}\pi r^3(t) - \mathcal{V}'_C} - C_\Sigma^o(t) \right]
 \end{aligned}$$

Osmotic Response Dynamics: RBC




$$\frac{d\mathcal{V}^i(t)}{dt} = A_0 RT\mathcal{L}_V\left(\frac{N_\Sigma^i}{\mathcal{V}^i(t)} - C_\Sigma^o(t)\right)$$

Primary osmotic responses of cells: Dynamics



Simple cases


- spherical cell (egg): $A(t) = 4\pi r^2(t); \quad \mathcal{V}^i(t) = \frac{4}{3}\pi r^3(t) - \mathcal{V}'_C(t)$
- cylindrical cell (muscle fiber): $A(t) = 2\pi r(t)L; \quad \mathcal{V}^i(t) = \pi r^2(t)L - \mathcal{V}'_C(t)$
- concave cell (RBC): $A(t) = A_0$



Extracellular medium was changed at $t = 0$ from 100% seawater to 60% seawater

Figure 4.28


Primary osmotic responses of cells: Dynamics



Simple cases

- spherical cell (egg): $A(t) = 4\pi r^2(t); \quad V^i(t) = \frac{4}{3}\pi r^3(t) - V_C^i(t)$
- cylindrical cell (muscle fiber): $A(t) = 2\pi r(t)L; \quad V^i(t) = \pi r^2(t)L - V_C^i(t)$
- concave cell (RBC): $A(t) = A_0$

→ Can simulate numerically



$$\alpha_v = \frac{RTL_V A^n C_\Sigma^{in}}{V^{in}}$$

Conclusions from Water Transport Experiments

- Both equilibrium and kinetic responses of simple cells (eggs) consistent with simple model of swelling based on macroscopic laws of osmosis
- However, there are some important issues that don't fit
 - water specialists: \mathcal{L}_V varies by factor of 1000 across cell types (RBC's are very permeant, artificial lipid bilayers are not)
 - Hg compounds: small amounts of Hg \rightarrow large $\Delta\mathcal{L}_V$ (little effect on artificial lipid bilayers)
 - hormonal control: vasopressin
 - antidiuretic hormone, secreted by hypothalamus
 - controls water retention in kidney
 - mechanism is to alter \mathcal{L}_V in water transporting epithelia

Molecular Aspects of Water Channels

Water Channels (Aquaporins)

"the plumbing system for cells"

First sequenced by Preston and Agre (1991)

- CHIP: CHannel-forming Integral Protein
- 807 base pairs in cDNA → 269 amino acids
(sequence provides 1D structure)
- hydrophobicity: 6 membrane spanning sections
(20 amino acids organized in alpha helix)
→ suggests 2D structure
- 4 proteins per water channel

→ 2003 Nobel Prize (Agre & MacKinnon)

Aquaporin




Figure 4.31

CHIP =
channel-forming integral protein




Figure 4.33

Aquaporin






Figure 1.37

Vasopressin

→ hormone that regulates water retention
(via affecting the membrane permeability)







Figure 2.19

Exercise

A volume element with constant cross-sectional area A has rigid walls and is divided into two parts by a rigid, semi-permeable membrane that is mounted on frictionless bearings so that the membrane is free to move in the x -direction as shown in the following figure.



The semi-permeable membrane is permeable to water but not to the solutes (glucose or NaCl or CaCl_2). At $t = 0$, solute 1 is added to side 1 to give an initial concentration of $c_1(0)$ and solute 2 is added to side 2 to give an initial concentration of $c_2(0)$. Concentrations are specified as the number of milli-moles of glucose or NaCl or CaCl_2 per liter of solution. The initial position of the membrane is $x(0)$. For the following values, find the final (equilibrium) values of the membrane position $x(\infty)$, and the concentrations, $c_1(\infty)$ and $c_2(\infty)$:


$$c_{glucose}^1(0) = 0; c_{glucose}^2(0) = 10; x(0) = 5.$$

$$c_{glucose}^1(0) = 30; c_{NaCl}^2(0) = 20; x(0) = 4.$$

The semi-permeable membrane is permeable to water but not to the solutes (glucose or NaCl or CaCl²). At $t = 0$, solute 1 is added to side 1 to give an initial concentration of $c_1(0)$ and solute 2 is added to side 2 to give an initial concentration of $c_2(0)$. Concentrations are specified as the number of milli-moles of glucose or NaCl or CaCl² per liter of solution. The initial position of the membrane is $x(0)$. For the following values, find the final (equilibrium) values of the membrane position $x(\infty)$, and the concentrations, $c_1(\infty)$ and $c_2(\infty)$:

$$c_{glucose}^1(0) = 0; c_{glucose}^2(0) = 10; x(0) = 5.$$

$$c_{glucose}^1(0) = 30; c_{NaCl}^2(0) = 20; x(0) = 4.$$



ANS: Keep in mind that the salt will disassociate. So when considering the osmolarity for each case:

- first scenario will cause the membrane to move all the way to the left ($c_1(\infty)$ will be undefined and $c_2(\infty)=5$)
- second scenario requires for osmotic equilibrium that $c_1(\infty) = 2c_2(\infty)$. Thus $c_1(\infty) = 36$, $c_2(\infty) = 18$, and $x(\infty) = 10/3$.

Exercise

A long and thin cylindrical cell of radius r and length l contains N_{Σ}^i moles of the impermeant solute S , and is immersed in a bath whose concentration of S is C_{Σ}^o . The membrane has a hydraulic conductivity \mathcal{L}_V . Assume that the membrane of the cell is permeable only to water and that the volume of water in the cell equals the volume of the cell. Also assume that flow of water through the ends of the cylindrical cell is negligible and that, as the cell changes its volume, only the radius changes while the length of the cell is constant. Assume that the hydraulic pressure difference across the membrane is zero.

a) Show that the cell radius r satisfies the differential equation

$$\frac{dr(t)}{dt} + \frac{A}{r^2(t)} = B.$$

b) Find A and B in terms of l , \mathcal{L}_V , N_{Σ}^i , C_{Σ}^o , R , and T , where R is the molar gas constant and T is absolute temperature.

ANS:

a. The equation of water volume conservation is

$$-\frac{1}{A(t)} \frac{d\mathcal{V}_c(t)}{dt} = \mathcal{L}_V RT \left(C_{\Sigma}^o - \frac{N_{\Sigma}^i}{\mathcal{V}_c(t)} \right).$$

If the surface areas of the circular ends of the cell are ignored, then the surface area of the cell is $A(t) = 2\pi r(t)l$ and its volume is $\mathcal{V}_c(t) = \pi r^2(t)l$. Substitution of these relations into the conservation relation yields

$$\begin{aligned} -\frac{1}{2\pi r(t)l} \frac{d(\pi r^2(t)l)}{dt} &= \mathcal{L}_V RT \left(C_{\Sigma}^o - \frac{N_{\Sigma}^i}{\pi r^2(t)l} \right), \\ -\frac{dr(t)}{dt} &= \mathcal{L}_V RT \left(C_{\Sigma}^o - \frac{N_{\Sigma}^i}{\pi r^2(t)l} \right), \\ \frac{dr(t)}{dt} - \frac{\mathcal{L}_V RT N_{\Sigma}^i / (\pi l)}{r^2(t)} &= -\mathcal{L}_V RT C_{\Sigma}^o. \end{aligned}$$

The last equation has the form

$$\frac{dr(t)}{dt} + \frac{A}{r^2(t)} = B.$$

b. The constants are

$$\begin{aligned} A &= -\frac{\mathcal{L}_V RT N_{\Sigma}^i}{\pi l}, \\ B &= -\mathcal{L}_V RT C_{\Sigma}^o. \end{aligned}$$

