











VSEPR Water



The oxygen is in the centre, the hydrogens are at two of the vertices. The lone pairs point to the other two. The molecule is bent.

Chemical bonding II 7

Chemical bonding II 8

CHEM 1000 3.0

Bond Angles

- This analysis suggests that all three molecules should have bond angles of 109.5°.
 - The methane bond angle is $109.5^{\rm o}$ but for ammonia it is $107^{\rm o}$ and in water it is $104.5^{\rm o}$
 - The lone pair electrons are not constrained as much as the bonding pairs. They spread out thus the repulsive forces are:
- Lone pair-lone pair>Lone pair>bond pair>bond pair

CHEM 1000 3.0

Molecular Geometry

- VSEPR theory can be used to describe the shapes of most molecules.
- Warnings
 - When you describe the shape, don't include the lone pairs. (water is bent, not tetrahedral)
 - Molecules have three dimensions (methane is a tetrahedron not a square)

CHEM 1000 3.0

• Be familiar with table 10.1 pg 399-400

TABLE 10.1 Molecular Geometry as a Function of Electron-Group Geometry							
Number of Electron Groups	Electron- Group Geometry	Number of Lone Pairs	VSEPR Notation	Molecular Geometry	Ideal Bond Angles	Example	VSEPR Basic Molecular Configurations models
2	linear	0	AX ₂	(linear)	180°	BeC12	0-0-0
⁸ For a discussion of the structure of SO ₂ , see page 402. ^b For a discussion of the placement of the lone-pair electrons in this structure, see page 401.							
Copyright © 2007 Pearson Prentice Hall, Inc.							
CHEM 1000 3.0							Chemical bonding II 10























Going Beyond Lewis Structures

- Lewis structures are very useful in explaining the bonding in simple molecules and in predicting molecular shapes.
- They do not explain why electrons in bond pairs bring nuclei together.
- They can not be used to estimate bond lengths or bond strengths.

CHEM 1000 3.0

Going Beyond Lewis Structures

- There are two currently used bonding theories, valence bond theory and molecular orbital theory.
- Valence bond theory envisions bonding as resulting from the overlap of atomic orbitals.
- Molecular orbital theory moves past atomic orbitals and derives orbitals that belong to the molecule as a whole.
- We will concentrate on valence bond theory as that is sufficient to rationalize the structures we see.

CHEM 1000 3.0

Chemical bonding II 17

Chemical bonding II 16

Valence Bond Method

- This method considers what happens to the valence orbitals (the outer shell).
- If the atoms start a long way apart, there is no interaction.
- As the atoms move closer together the orbitals may overlap.
- If there is an electron in the orbital then there is a high probability of finding an electron between the nuclei.
- This is the region where there is a covalent bond.

CHEM 1000 3.0 Chemical bonding II 18

















Valence Bond Rethinking

- We can no longer assume that the orbitals in a bonded atom are the same as those of an isolated atom.
- Remember that the atomic orbitals come from solving an equation assuming a single nucleus. In a molecule the electron has to respond to more nuclei.
- We should solve the "new" problem, but we usually say we can combine the atomic orbitals in some way.

CHEM 1000 3.0

Valence Bond method for methane

- Since the orbitals are wave functions, we can take the 2s and 2p wave functions and combine them to give 4 equivalent wave functions. These can be made to have the same shape and energy.
- This is called hybridization and the resulting orbitals are called hydrid orbitals.
- The combination of one "s" and three "p" orbitals gives four "sp³" hybrid orbitals.

CHEM 1000 3.0

Chemical bonding II 26





















- NH_3 has sp³ bonding and a bond angle of ~109.
- PH₃ does not hybridize.
- WHY NOT?
- The energy gain in forming the bonds is not enough to get back the energy to make the hybrids.
- This is because the energy separation between 3s and 3p is higher than 2s and 2p on a relative basis.
- Going down a main group in the periodic table, elements tend to make less use of valence s electrons and form weaker bonds with smaller angles.

CHEM 1000 3.0

• The sp³ hybrid orbitals don't explain the bonding for Boron (group 13) or for Beryllium (group 2)



• The problem appears to be that there are insufficient electrons to even half fill the 4 sp³ orbitals.

- In these case we need to consider different hybridization schemes.
- We define hybrids such that the number of hybrids is the same as the number of valence electrons.

CHEM 1000 3.0







































• The overlap of the "p" orbitals can be thought of as giving the three double bonds, but in reality the π orbital system is delocalized around the whole ring. This eliminates the need to discuss resonance.

CHEM 1000 3.0

Other kinds of "bonds"

- In the section on gases we said there were forces between the molecules. Now we know more about bonding and electronic structure we can revisit the question.
- What kinds of forces exist between molecules?

CHEM 1000 3.0 Chemical bonding II 49

Van der Waal Forces

- Van der Waal forces are those attractive forces between molecules that are responsible for:
 - Real gas behaviour (Van der Waal "a")Condensation
- There are a number of such forces resulting in different types of interactions between molecules.

CHEM 1000 3.0

Chemical bonding II 50

Van der Waal Forces London or Dispersion Force

- The quantum concept of electronic structure talks about the probability of an electron being in a certain region at a certain time.
- Even for an atom or a perfectly symmetrical molecule there is a finite chance that the electronic charges are not uniformly distributed.
- This produces an instantaneous dipole.

CHEM 1000 3.0 Chemical bonding II 51





Van der Waal Forces London or Dispersion Force

- This force is going to be strongest in a molecule with a large number of electrons or for an elongated molecule.
- The ability or tendency of a molecule to have charge separation occur is called polarizability.
- The effect of London forces can often be seen in the boiling points of similar compounds.

CHEM 1000 3.0









• How do we tell if a molecule has a dipole?

We can put the molecules in an electric field and see the effects of them "lining up".





- Not all molecules with polar bonds have a dipole.
- This is because the dipoles can cancel each other out.

CHEM 1000 3.0

Chemical bonding II 58



Van der Waal Forces Dipole-Dipole interactions

CHEM 1000 3.0

Chemical bonding II 60

•20

• For molecules with permanent dipoles, Dipole-Dipole interactions can occur.

Van der Waal Forces Dipole-Dipole interactions

- Dipole-dipole interactions are in addition to the London forces.
- The overall interaction between polar molecules is higher than for nonpolar molecules.



Chemical bonding II 61

Chemical bonding II 62

CHEM 1000 3.0



Hydrogen Bonding

- This is an extreme case.
- Hydrogen atoms bonded to very highly electronegative atoms can form weak bonds with the atom in an adjacent molecule.
- This only occurs for hydrogen bonded to F, O, and N.
- The bond energy is 15 40 kJ mol⁻¹, high compared to London forces but low compared to covalent bonds (>150 kJ mol⁻¹)

CHEM 1000 3.0

Effect of Hydrogen Bonding 400 Normal boiling point, K 200 -100 -H₂Te H₂Se SbH₂ AsH₃ HBr HI SnH₄ PH HCI GeH₄ SiH CH₄ 0 | 75 125 150 50 100 25 Molecular mass, u Chemical bonding II 63 CHEM 1000 3.0









