

Thermochemistry

- Petrucci, Harwood and Herring: Chapter 7

CHEM 1000 3.0

Thermochemistry 1

Thermochemistry

- The study of energy in chemical reactions
- A sub-discipline of thermodynamics
- Thermodynamics studies the bulk properties of matter and deduces a few general laws
 - It does not require any knowledge/assumptions of molecules

CHEM 1000 3.0

Thermochemistry 2

Definitions

- System
 - The part of the universe we chose to study
- Surroundings
 - The rest of the universe (normally we only worry about the immediate surroundings)
- Process
 - A physical occurrence (usually involving energy flow)

CHEM 1000 3.0


Thermochemistry 3

Additional definitions

- Open system
 - A system where energy and matter can be exchanged with the surroundings
- Closed system
 - A system where energy but not matter can be exchanged with the surroundings
- Isolated system
 - A system where neither energy nor matter can be exchanged with the surroundings

CHEM 1000 3.0

Thermochemistry 4



CHEM 1000 3.0

Thermochemistry 5

Systems and Energy

- All systems will contain energy
 - In thermodynamics we are interested in the flow of energy, particularly in the forms of heat and work.
 - Note that heat and work occur when there is a process. They only exist when something happens.
 - The system has energy, (often described as the capacity to do work), it does not have heat or work.

CHEM 1000 3.0

Thermochemistry 6

Heat

- Heat

- Energy that is transferred between a system and its surroundings as a result of temperature differences
- Heat transfer can change the temperature of something but it does not always do (only) that
- Heat transfer can melt or vaporize material

CHEM 1000 3.0

Thermochemistry 7

- In the case of a material that does not change phase, the increase in temperature of a system ΔT due to the input of a given amount of heat q is given by

$$q = c \Delta T$$

Diagram showing the components of the equation $q = c \Delta T$:

- q is labeled as heat (J)
- c is labeled as heat capacity (J K^{-1})
- ΔT is labeled as temperature change (K)

CHEM 1000 3.0

Thermochemistry 8

- The heat capacity is a constant that depends on the system. So it's not particularly useful.
- It is better to be able to define heat capacity in terms of a particular compound

$$q = nC\Delta T$$

Diagram showing the components of the equation $q = nC\Delta T$:

- n is labeled as Number of moles
- C is labeled as molar heat capacity ($\text{J K}^{-1} \text{mol}^{-1}$)

$$q = mc\Delta T$$

Diagram showing the components of the equation $q = mc\Delta T$:

- m is labeled as mass (kg)
- c is labeled as specific heat ($\text{J K}^{-1} \text{kg}^{-1}$)

CHEM 1000 3.0

Thermochemistry 9

Units

- The SI unit of heat is a Joule (since it is an energy)
- The older unit of heat is the calorie which is defined as the heat required to raise the temperature of 1 g of water 1°C
 $1\text{ cal} = 4.184\text{ J}$

CHEM 1000 3.0

Thermochemistry 10

Sign convention and conservation of energy

- When heat flows between a system and its surroundings, we define:
q to be positive if heat is supplied to the system
q to be negative if heat is withdrawn from the system
- If there are no phase changes, conservation of energy requires that

$$q_{\text{system}} + q_{\text{surroundings}} = 0$$

or

$$q_{\text{system}} = -q_{\text{surroundings}}$$

CHEM 1000 3.0

Thermochemistry 11

Work

- Often when a chemical reaction occurs, work is done. (this is the principle of an engine)
- Since work and heat are both forms of energy we must consider both.

CHEM 1000 3.0

Thermochemistry 12

Work

- Work is done when a force acts through a distance. For example when a mass is moved.
- Definition

$$w = F \times d$$

Work (J) Force (N) Distance (m)

CHEM 1000 3.0

Thermochemistry 13

- The same definitions apply for work as for heat:

w is positive if work is done on the system

w is negative if work is done by the system


CHEM 1000 3.0

Thermochemistry 14

Pressure-Volume Work

This is the most common type of work.

For example, a gas formed in a reaction pushes against the atmosphere causing a volume change.




(Note that because the gas is expanding, the system is doing work on the surroundings, so work must be negative.)

CHEM 1000 3.0

Thermochemistry 15

Pressure-Volume Work



$$\begin{aligned}
 w &= -F \times d \\
 &= -(M \times g) \times \Delta h \\
 &= \frac{-(M \times g)}{A} \times \Delta h \times A \\
 &= -P\Delta V
 \end{aligned}$$

The gas is expanding in this process so the system is doing work on the surroundings and therefore it is negative.

Note: Derivation assumes that the piston and pan assembly is massless and the water bath keeps the temperature of the gas constant.

CHEM 1000 3.0

Thermochemistry 16

Calculating Pressure-Volume Work

Suppose the gas in the previous figure is 0.100 mol He at 298 K and the each mass in the figure corresponds to an external pressure of 1.20 atm. How much work, in Joules, is associated with its expansion at constant pressure?

Assume an ideal gas and calculate the volume change:

$$\begin{aligned}
 V_i &= nRT/P \\
 &= (0.100 \text{ mol})(0.08201 \text{ L atm mol}^{-1} \text{ K}^{-1})(298 \text{ K})/(2.40 \text{ atm}) \\
 &= 1.02 \text{ L} \\
 V_f &= 2.04 \text{ L} \qquad \Delta V = 2.04 \text{ L} - 1.02 \text{ L} = 1.02 \text{ L}
 \end{aligned}$$

CHEM 1000 3.0

Thermochemistry 17

Calculate the work done by the system:

$$\begin{aligned}
 w &= -P\Delta V \\
 &= -(1.20 \text{ atm})(1.02 \text{ L}) \left(\frac{101 \text{ J}}{1 \text{ L atm}} \right) \\
 &= -1.24 \times 10^2 \text{ J}
 \end{aligned}$$

A negative value signifies that work is done ON the surroundings

Where did the conversion factor come from?

Compare two versions of the gas constant and calculate.

$$8.3145 \text{ J/mol K} = 0.082057 \text{ L atm/mol K}$$


$$(8.3145 \text{ J/mol K}) / (0.082057 \text{ L atm/mol K}) = 101.33 \text{ J/L atm}$$

CHEM 1000 3.0

Thermochemistry 18

First Law of Thermodynamics

- We have defined a system, work and heat.
- A closed system can exchange heat and work with its surroundings.
- The system has some energy we call the internal energy, U.



CHEM 1000 3.0

Thermochemistry 19

First Law of Thermodynamics

- The first law of thermodynamics

$$\Delta U = q + w$$

CHEM 1000 3.0

Thermochemistry 20

The internal energy is a **state function**. This means that its value depends on the state of the system (its pressure, temperature etc) not on how it got there.

Heat and work are not state functions. Their values depend on where the system has been, not on where it is.

CHEM 1000 3.0

Thermochemistry 21

Where is the internal energy?

- It is not necessary that we know where the internal energy resides, But:
- It can be energy due to motion in the molecules e.g. translational energy (kinetic), or energy due to molecular rotational and vibration.
- For an ideal gas the internal energy is the kinetic energy

$$E = \frac{3RT}{2}$$

CHEM 1000 3.0

Thermochemistry 22

Thermochemistry and the First Law

We have $\Delta U = q + w$

(For chemical reactions, the most common form of heat is the heat of reaction q_{rxn} .)

For a constant volume process:

$$\Delta U = q + w = q - P \Delta V$$

But $\Delta V = 0$ therefore $\Delta U = q_v$


The subscript indicates a constant volume process.

CHEM 1000 3.0

Thermochemistry 23

More reactions occur at constant pressure conditions than constant volume. In this case the volume of the system can change and work can be done.

Suppose the system has to change from one state to another. It can do it at constant volume or constant pressure but the heat involved is different



CHEM 1000 3.0

Thermochemistry 24

Internal energy

Initial state U_i

Final state U_f

(a) $\Delta U = q_v$

(b) $\Delta U = q_p + w$

$q_v = q_p + w$

But $q_v = \Delta U$ and $w = -P \Delta V$

So

$\Delta U = q_p - P \Delta V$

$q_p = \Delta U + P \Delta V$

U, P and V are state functions so a combination of them will also be a state function.

Define **enthalpy** as: $H = U + PV$

$\Delta H = \Delta U + \Delta PV$

At constant pressure

$\Delta H = \Delta U + P \Delta V$ so $\Delta H = q_p$

CHEM 1000 3.0 Thermochemistry 25

$2 \text{CO(g)} + \text{O}_2\text{(g)} \rightarrow 2 \text{CO}_2\text{(g)}$

At constant volume

$\Delta U = q_v = -563.5 \text{ kJ}$

At constant pressure, work is done on the system. You get the heat from reaction and from the work.

$\Delta H = q_p = -566.0 \text{ kJ}$

CHEM 1000 3.0 Thermochemistry 26

Enthalpy and change of state

- There is usually a flow of heat when there is a change of state (solid to liquid, liquid to vapour)
- This is usually expressed as the enthalpy of fusion or enthalpy of vaporization. E.g.


$$\text{H}_2\text{O(s)} \rightarrow \text{H}_2\text{O(l)} \quad \Delta H = 6.01 \text{ kJ mol}^{-1}$$

$$\text{H}_2\text{O(l)} \rightarrow \text{H}_2\text{O(g)} \quad \Delta H = 44.0 \text{ kJ mol}^{-1}$$
- This is often just called the heat of fusion or heat of vaporization.

CHEM 1000 3.0 Thermochemistry 27


Enthalpy/thermochemistry

- Thermochemistry is the branch of thermodynamics that deals with energy changes in chemical reactions.
- Normally a chemical reaction is expressed with the associated enthalpy stated



CHEM 1000 3.0

Thermochemistry 28



CHEM 1000 3.0

Thermochemistry 29


Hess's Law


Enthalpy is a state function. Therefore we can use some obvious properties to determine unknown enthalpies.

- Enthalpy changes are proportional to the amount of material.
- The sign of the enthalpy changes if a reaction is reversed.
- (Hess's Law) If a reaction can be thought of as proceeding through a number of steps, then the enthalpy change on reaction must be the sum of the enthalpy changes for each step.

CHEM 1000 3.0

Thermochemistry 30





Standard enthalpies of formation

- We have seen that the enthalpy of a reaction is calculated as a difference between the enthalpies of the products and reactants.
- We do not have an absolute scale for enthalpies (what is the enthalpy of O_2 ? For example)
- Just as we measure heights above sea-level, we measure enthalpies with respect to the standard state.

CHEM 1000 3.0 Thermochemistry 33

Standard State

- By definition: the enthalpy of an element in its most stable form at a pressure of 1 bar for a specified temperature is zero. (usually tabulated for 298K)
- E.g. $\text{O}_2(\text{g})$, $\text{Br}_2(\text{l})$, $\text{C}(\text{graphite})$, $\text{Na}(\text{s})$...
- We calculate all enthalpies from this standard state

CHEM 1000 3.0


Thermochemistry 34

- We can now define a standard enthalpy of formation as the enthalpy change that occurs in the formation of one mole of the substance from its elements in their standard states.
- Usually given the symbol ΔH_f° where the f is for formation and the $^\circ$ indicates standard conditions.

CHEM 1000 3.0

Thermochemistry 35

- Examples of standard enthalpies of formation



CHEM 1000 3.0

Thermochemistry 36

- Now we have standard enthalpies of formation we can determine the enthalpy of a reaction.

$$\Delta H_{\text{reaction}} = H_{\text{products}} - H_{\text{reactants}}$$

$$= \sum v_p \Delta H_f^\circ(\text{products}) - \sum v_r \Delta H_f^\circ(\text{reactants})$$


where Σ is a sum over the standard enthalpies of formation ΔH_f° multiplied by their stoichiometric coefficients v .

CHEM 1000 3.0

Thermochemistry 37

Where do ΔH_f° values come from?


- All enthalpies of formation can be traced to calorimetry experiments.
- Most widely used is the bomb calorimeter



CHEM 1000 3.0

Thermochemistry 38

- The bomb calorimeter is a constant volume reactor, usually used for combustion reactions.
- The reagents are the system and the calorimeter is the surroundings.
- By measuring the heat transferred into the surroundings the heat of the reaction can be determined.



CHEM 1000 3.0

Thermochemistry 39

- If the increase in temperature of the surroundings is ΔT then the heat going into the surroundings is

$$q_{\text{calorimeter}} = c \Delta T$$

Since the reaction takes place at constant volume

$$\Delta U = q_{\text{rxn}} = -q_{\text{calorimeter}} \quad (\text{converted to per mole})$$

To get ΔH_{rxn} we use

$$\Delta H_{\text{rxn}} = \Delta U + \Delta PV$$

And the ideal gas law

$$\Delta H_{\text{rxn}} = \Delta U + \Delta nRT$$

Then we use

$$\Delta H_{\text{rxn}} = \sum_p \nu_p \Delta H_f^\circ(\text{products}) - \sum_r \nu_r \Delta H_f^\circ(\text{reactants})$$


CHEM 1000 3.0

Thermochemistry 40

Example:

Calculate the standard heat of formation of CH_4 if a bomb calorimeter experiment at an initial temperature of 298K gives a measured heat gain in the calorimeter of 885.4 kJ per mole of CH_4 .

The combustion reaction is the one that fully oxidizes the reagents:



$$\Delta U = q_{\text{rxn}} = -q_{\text{calorimeter}} = -885.4 \text{ kJ mol}^{-1}$$


$$\Delta H_{\text{rxn}} = \Delta U + \Delta nRT$$

$$\Delta n = 1 - 3 = -2 \quad \Delta H_{\text{rxn}} = -885.4 + (-2) \times 8.314 \times 298 \times 10^{-3} \text{ kJ mol}^{-1} = -890.4 \text{ kJ mol}^{-1}$$

$$\Delta H_{\text{rxn}} = \sum_p \nu_p \Delta H_f^\circ(\text{products}) - \sum_r \nu_r \Delta H_f^\circ(\text{reactants})$$

CHEM 1000 3.0

Thermochemistry 41



$$\Delta H_{\text{rxn}} = \sum_p \nu_p \Delta H_f^\circ(\text{products}) - \sum_r \nu_r \Delta H_f^\circ(\text{reactants})$$

$$\Delta H_{\text{rxn}} = \Delta H_f^\circ(\text{CO}_2) + 2\Delta H_f^\circ(\text{H}_2\text{O}) - \Delta H_f^\circ(\text{CH}_4)$$

If we know the heats of formation of CO_2 and H_2O then:

$$\Delta H_{\text{rxn}} = \Delta H_f^\circ(\text{CO}_2) + 2\Delta H_f^\circ(\text{H}_2\text{O}) - \Delta H_f^\circ(\text{CH}_4)$$

$$-890.4 = -393.51 + 2 \times (-285.85) - \Delta H_f^\circ(\text{CH}_4)$$

$$\Delta H_f^\circ(\text{CH}_4) = -74.8 \text{ kJ mol}^{-1}$$

CHEM 1000 3.0




Thermochemistry 42

Bond Energies

- The Bond Dissociation Energy is the energy to break a mole of bonds.



- For cases where there are more than one identical bond, the average is taken



CHEM 1000 3.0

Thermochemistry 43

Bond Energies

Bond	Bond Energy, kJ/mol	Bond	Bond Energy, kJ/mol	Bond	Bond Energy, kJ/mol
H—H	436	C—C	347	N—N	163
H—C	414	C=C	611	N=N	418
H—N	389	C≡C	837	N≡N	946
H—O	464	C—N	305	N—O	222
H—S	368	C≡N	615	N=O	590
H—F	565	C=N	891	O—O	142
H—Cl	431	C—O	360	O=O	498
H—Br	364	C=O	736 ^b	F—F	159
H—I	297	C—Cl	339	Cl—Cl	243
				Br—Br	193
				I—I	151

* Although all data are listed with about the same precision (three significant figures), some values are actually known more precisely. Specifically, the values for the diatomic molecules H_2 , HF, HCl, HBr, HI, N_2 ($\text{N}=\text{N}$), O_2 ($\text{O}=\text{O}$), F_2 , Cl_2 , Br_2 , and I_2 are actually bond-dissociation energies, rather than average bond energies.

^b The value for the C=O bonds in CO_2 is 799 kJ/mol.

Copyright © 2007 Pearson Prentice Hall, Inc.

CHEM 1000 3.0

Thermochemistry 44

Bond Energies

- Hess's law can be used to show

$$\Delta\text{H}_{\text{rxn}} = \Delta\text{H}(\text{bond breakage}) - \Delta\text{H}(\text{bond formation})$$

$$\Delta\text{H}_{\text{rxn}} = \sum \text{BE}(\text{reactants}) - \sum \text{BE}(\text{products})$$

This will never be as precise as using heats of formation as BEs are always averages.


CHEM 1000 3.0

Thermochemistry 45

Bond Energy Examples

1) Example 10-15 pg 409

Calculate ΔH_{rxn} for the reaction



- draw each molecule and its bonds to emphasize which bonds are broken and which ones are formed (could break all bonds in reactants and then form all of the bonds in the products, but this is not necessary)
- one C-H bond and one Cl-Cl bond are broken; one C-Cl bond and one H-Cl bond are formed


$$\begin{aligned}\Delta H_{\text{rxn}} &= \Delta H(\text{bonds broken}) + \Delta H(\text{bonds formed}) \\ &= \{\text{BE}(\text{C-H}) + \text{BE}(\text{Cl-Cl})\} - \{\text{BE}(\text{C-Cl}) + \text{BE}(\text{H-Cl})\} \\ &= \{414 \text{ kJ} + 243 \text{ kJ}\} - \{339 \text{ kJ} + 431 \text{ kJ}\} \\ &= -113 \text{ kJ}\end{aligned}$$

CHEM 1000 3.0

Thermochemistry 46

2) Practice Problem - estimate versus experiment

Calculate ΔH_{rxn} for the reaction



- note that it is important to know the bond order
- one N≡N triple bond and 3 H-H single bonds are broken; 6 N-H single bonds are formed (sketch if necessary)

$$\begin{aligned}\Delta H_{\text{rxn}} &= \Delta H(\text{bonds broken}) + \Delta H(\text{bonds formed}) \\ &= \{\text{BE}(\text{N}\equiv\text{N}) + 3 \text{BE}(\text{H-H})\} - \{6 \text{BE}(\text{N-H})\} \\ &= +946 \text{ kJ} + 3(436 \text{ kJ}) - 6(389 \text{ kJ}) \\ &= -80 \text{ kJ}\end{aligned}$$

Note: experimentally ΔH_{rxn} determined to be -92 kJ

CHEM 1000 3.0

Thermochemistry 47
