

7


Newton's Third Law

7.1 Interacting Objects


7.2 Analyzing Interacting Objects

Exercises 1–7: Follow steps 1–3 of Tactics Box 7.1 to draw interaction diagrams describing the following situations. Your diagrams should be similar to Figures 7.6 and 7.10.


1. A bat hits a ball.



2. A massless string pulls a box across the floor. Friction is not negligible.



3. A boy pulls a wagon by a rope attached to the front of the wagon. The rope is not massless, and rolling friction is not negligible.




4. A skateboarder is pushing on the ground to speed up. Treat the person and the skateboard as separate objects.

5. The bottom block is pulled by a massless string. Friction is not negligible. Treat the two blocks as separate objects.




6. A crate in the back of a truck does not slip as the truck accelerates forward. Treat the crate and the truck as separate objects.

7. The bottom block is pulled by a massless string. Friction is not negligible. Treat the pulley as a separate object.



7.3 Newton's Third Law

8. Block A is pushed across a horizontal surface at a *constant* speed by a hand that exerts force $\vec{F}_{H \text{ on } A}$. The surface has friction.



a. Draw two free-body diagrams, one for the hand and the other for the block. On these diagrams:

- Show only the *horizontal* forces, such as was done in Figure 7.14 of the text.
- Label force vectors, using the form $\vec{F}_{C \text{ on } D}$.
- Connect action/reaction pairs with dotted lines.
- On the hand diagram show only $\vec{F}_{A \text{ on } H}$. Don't include $\vec{F}_{\text{body on } H}$.
- Make sure vector lengths correctly portray the relative magnitudes of the forces.


b. Rank in order, from largest to smallest, the magnitudes of *all* of the horizontal forces you showed in part a. For example, if $F_{C \text{ on } D}$ is the largest of three forces while $F_{D \text{ on } C}$ and $F_{D \text{ on } E}$ are smaller but equal, you can record this as $F_{C \text{ on } D} > F_{D \text{ on } C} = F_{D \text{ on } E}$.

Order:

Explanation:

c. Repeat both part a and part b for the case that the block is *speeding up*.

9. A second block B is placed in front of Block A of question 8. B is more massive than A: $m_B > m_A$. The blocks are speeding up.



- a. Consider a *frictionless* surface. Draw *separate* free-body diagrams for A, B, and the hand. Show only the horizontal forces. Label forces in the form $\vec{F}_{C \text{ on } D}$. Use dashed lines to connect action/reaction pairs.


- b. By applying the second law to each block and the third law to each action/reaction pair, rank in order *all* of the horizontal forces, from largest to smallest.

Order:

Explanation:

- c. Repeat parts a and b if the surface has friction. Assume that A and B have the same coefficient of kinetic friction.

10. Blocks A and B are held on the palm of your outstretched hand as you lift them straight up at *constant speed*. Assume $m_B > m_A$ and that $m_{\text{hand}} = 0$.



- a. Draw *separate* free-body diagrams for A, B, and your hand.
- Show *all* vertical forces, including the gravitational forces on the blocks.
 - Make sure vector lengths indicate the relative sizes of the forces.
 - Label forces in the form $\vec{F}_{C \text{ on } D}$
 - Connect action/reaction pairs with dashed lines.


b. Rank in order, from largest to smallest, all of the vertical forces. Explain your reasoning.

11. A mosquito collides head-on with a car traveling 60 mph.


- a. How do you think the size of the force that the car exerts on the mosquito compares to the size of the force that the mosquito exerts on the car?

- b. Draw *separate* free-body diagrams of the car and the mosquito at the moment of collision, showing only the horizontal forces. Label forces in the form $\vec{F}_{C \text{ on } D}$. Connect action/reaction pairs with dotted lines.


Exercises 12–16: Write the acceleration constraint in terms of *components*. For example, write $(a_1)_x = (a_2)_y$, if that is the appropriate answer, rather than $\vec{a}_1 = \vec{a}_2$.




Constraint: _____




Constraint: _____



Constraint: _____



Constraint: _____




Constraint: _____


7.4 Ropes and Pulleys

Exercises 17–22: Determine the reading of the spring scale.


- All the masses are at rest.
- The strings and pulleys are massless, and the pulleys are frictionless.
- The spring scale reads in kg.




Scale = _____




Scale = _____




Scale = _____



Scale = _____




Scale = _____



Scale = _____

7.5 Examples of Interacting-Objects Problems

23. Blocks A and B, with $m_B > m_A$, are connected by a string. A hand pushing on the back of A accelerates them along a frictionless surface. The string (S) is massless.



- a. Draw separate free-body diagrams for A, S, and B, showing only horizontal forces. Be sure vector lengths indicate the relative size of the force. Connect any action/reaction pairs with dotted lines.

- b. Rank in order, from largest to smallest, all of the horizontal forces. Explain.

- c. Repeat parts a and b if the string has mass.

- d. You might expect to find $F_{S \text{ on } B} > F_{H \text{ on } A}$ because $m_B > m_A$. Did you? Explain why $F_{S \text{ on } B} > F_{H \text{ on } A}$ is or is not a correct statement.


24. Blocks A and B are connected by a massless string over a massless, frictionless pulley. The blocks have just this instant been released from rest.
- a. Will the blocks accelerate? If so, in which directions?

- b. Draw a separate free-body diagram for each block. Be sure vector lengths indicate the relative size of the force. Connect any action/reaction pairs or "as if" pairs with dashed lines.


- c. Rank in order, from largest to smallest, all of the vertical forces. Explain.

- d. Compare the magnitude of the *net* force on A with the *net* force on B. Are they equal, or is one larger than the other? Explain.

- e. Consider the block that falls. Is the magnitude of its acceleration less than, greater than, or equal to g ? Explain.



25. In case a, block A is accelerated across a frictionless table by a hanging 10 N weight (1.02 kg). In case b, the same block is accelerated by a steady 10 N tension in the string.




Is block A's acceleration in case b greater than, less than, or equal to its acceleration in case a? Explain.

Blank space for the student's explanation.


Exercises 26–27: Draw separate free-body diagrams for blocks A and B. Connect any action/reaction pairs (or forces that act *as if* they are action/reaction pairs) together with dashed lines.

26.



Blank space for drawing free-body diagrams for exercise 26.

27.



Blank space for drawing free-body diagrams for exercise 27.